

GenCore version 4.5  
Copyright (c) 1993 - 2000 CompuGen Ltd.

OM protein - protein search, using sw model

Run on: August 13, 2001, 12:22:04 ; Search time 59.73 Seconds  
(without alignments)  
534.888 Million cell updates/sec

Title: US-09-784-340-2

Perfect score: 2802  
Sequence: 1 MRSKSAVFLDLQFCVGC.....KCFLECKKFKTKREKRE 527

Scoring table:  
BLOSUM62  
Gapop 10.0 , Gapept 0.5

Searched: 412676 seqs, 60623988 residues

Total number of hits satisfying chosen parameters: 412676

Minimum DB seq length: 0  
Maximum DB seq length: 200000000

Post-processing: Minimum Match 08  
Maximum Match 1008  
Listing first 45 summaries

Database :

A.Geneseq.0601:\*

1: /SIDSL/gcgdata/geneseq/geneseq/AA1980.DAT:\*  
2: /SIDSL/gcgdata/geneseq/geneseq/AA1981.DAT:\*  
3: /SIDSL/gcgdata/geneseq/geneseq/AA1982.DAT:\*  
4: /SIDSL/gcgdata/geneseq/geneseq/AA1983.DAT:\*  
5: /SIDSL/gcgdata/geneseq/geneseq/AA1984.DAT:\*  
6: /SIDSL/gcgdata/geneseq/geneseq/AA1985.DAT:\*  
7: /SIDSL/gcgdata/geneseq/geneseq/AA1986.DAT:\*  
8: /SIDSL/gcgdata/geneseq/geneseq/AA1987.DAT:\*  
9: /SIDSL/gcgdata/geneseq/geneseq/AA1988.DAT:\*  
10: /SIDSL/gcgdata/geneseq/geneseq/AA1989.DAT:\*  
11: /SIDSL/gcgdata/geneseq/geneseq/AA1990.DAT:\*  
12: /SIDSL/gcgdata/geneseq/geneseq/AA1991.DAT:\*  
13: /SIDSL/gcgdata/geneseq/geneseq/AA1992.DAT:\*  
14: /SIDSL/gcgdata/geneseq/geneseq/AA1993.DAT:\*  
15: /SIDSL/gcgdata/geneseq/geneseq/AA1994.DAT:\*  
16: /SIDSL/gcgdata/geneseq/geneseq/AA1995.DAT:\*  
17: /SIDSL/gcgdata/geneseq/geneseq/AA1996.DAT:\*  
18: /SIDSL/gcgdata/geneseq/geneseq/AA1997.DAT:\*  
19: /SIDSL/gcgdata/geneseq/geneseq/AA1998.DAT:\*  
20: /SIDSL/gcgdata/geneseq/geneseq/AA1999.DAT:\*  
21: /SIDSL/gcgdata/geneseq/geneseq/AA2000.DAT:\*  
22: /SIDSL/gcgdata/geneseq/geneseq/AA2001.DAT:\*

Pred. No. is the number of results predicted by chance to have a score greater than or equal to the score of the result being printed, and is derived by analysis of the total score distribution.

# SUMMARIES

Result No.	Score	Query Match	Length	ID	Description
1	1924.5	68.7	529	21	AA28677
2	1732	61.8	530	19	AA47126
3	1729	61.7	528	21	AA78933
4	1713	61.1	530	21	AA78935
5	1673.5	59.7	524	21	AA78934
6	1120	40.0	533	13	AA26153
7	1058.5	37.8	534	13	AA26154
8	755.5	27.0	245	21	AA57100
9	714.5	25.5	523	21	AA24025
10	714.5	25.5	523	21	AA59419
11	714.5	25.5	523	22	AA66168

12	708.5	25.3	523	22	AA68348	Human membrane or
13	405	14.5	78	21	AA603280	Human secreted pro
14	391	14.0	129	20	AA29525	Human lung tumour
15	391	14.0	129	21	AA44411	Human lung tumour
16	383.5	13.7	288	21	AA57092	UDP-glucuronosyltr
17	349.5	12.5	310	21	AA57098	UDP-glucuronosyltr
18	341	12.2	287	13	AA57096	UDP-glucuronosyltr
19	341	12.2	287	21	AA57096	UDP-glucuronosyltr
20	327.5	11.7	317	21	AA57097	UDP-glucuronosyltr
21	326	11.6	289	13	AA57093	UDP-glucuronosyltr
22	326	11.6	289	21	AA57094	UDP-glucuronosyltr
23	321	11.5	289	21	AA57095	UDP-glucuronosyltr
24	320	11.4	289	21	AA57095	UDP-glucuronosyltr
25	317.5	11.3	253	21	AA57099	UDP-glucuronosyltr
26	317	11.3	253	21	AA57099	UDP-glucuronosyltr
27	316	11.3	253	21	AA57099	UDP-glucuronosyltr
28	297.5	10.6	98	13	AA57096	UDP-glucuronosyltr
29	294.5	10.5	248	13	AA57096	UDP-glucuronosyltr
30	274.5	9.8	515	19	AA56750	Ecdysteroid UDP-gl
31	262	9.4	243	13	AA56750	Ecdysteroid UDP-gl
32	260	9.3	94	21	AA53721	Human colon cancer
33	254.5	9.1	506	12	AA53721	Ecdysteroid UDP-gl
34	235	8.4	74	13	AA530165	UGT1 Exon 4 produc
35	190	6.8	466	18	AA509825	Ecdysteroid UDP-gl
36	175	6.2	399	12	AA53989	UDP-glucose:thiohy
37	175	6.2	431	11	AA507464	zeaxanthin glycosy
38	175	6.2	431	20	AA507464	polyptide with e
39	171.5	6.1	68	21	AA56504	Protein encoded by
40	171.5	6.1	439	21	AA56504	Human prostate can
41	171.5	6.1	445	21	AA564893	Arabidopsis thalia
42	171.5	6.1	460	21	AA564892	Arabidopsis thalia
43	171.5	6.1	465	21	AA5640010	Arabidopsis thalia
44	171.5	6.1	460	21	AA564891	Arabidopsis thalia
45	171.5	6.1	481	21	AA5640009	Arabidopsis thalia

## ALIGNMENTS

RESULT	ID	Protein	Protein ID	Protein Description
1	AA28677	standard; Protein: 529 AA.		
XX	AA28677			
XX	AA28677			
DT	13-FEB-2001	(first entry)		
XX				
DE		Human carbohydrate-modifying enzyme Incyte ID No: 2912310CD1.		
XX				
KW		Human: carbohydrate-modifying enzyme: CME: antidiabetic:		
KW		Immunosuppressive; anti-HIV; antiinflammatory; antinaemic:		
KW		antiallergic; antihypertensive; antihypertensive; antihypertensive:		
KW		neurotrophic; antitumor; antitumor; antitumor; antitumor:		
KW		antiarthritic; antipsoriatic; uropathic; ophthalmologic:		
KW		dermatological; antitumor; cytotoxic; vincetristine; antidiabetic:		
KW		fungicide; protozoicide; tranquiliser; vulvar; diabetes:		
KW		autoimmune disorder; inflammatory disorder; infection.		
XX				
OS		Homo sapiens.		
PN				
XX				
PN		WO200063351-A2.		
XX				
PD		26-OCT-2000.		
XX				
XX		20-APR-2000; 2000WO-US10882.		
PF				
XX				
XX		21-APR-1999; 99US-0130383.		
PR				
XX				
XX				
PA		(INCYTE) INCYTE GENOMICS INC.		
XX				
XX				
PI		Ial P, Yue H, Tang YT, Hillman JL, Baughn MR, Yang J:		
XX				
DR		WPI; 2000-672729/65.		

DR N-PSDB: AAC65396.

XX Novel carbohydrate modifying enzyme polypeptides and polynucleotides  
PT for diagnosis, treatment, and prevention of carbohydrate metabolism  
PT disorders, autoimmune/inflammatory disorders, and cancer

PS Claim 1: Page 71-72; 75pp; English.

XX The present sequence is a human carbohydrate-modifying enzyme  
CC (CME). CME polynucleotides and polypeptides are useful for treating and  
CC diagnosing diseases associated with CME such as diabetes,  
CC autoimmune/inflammatory disorders such as AIDS, Addison's disease,  
CC adult respiratory distress syndrome, allergies, anaemia, asthma,  
CC atherosclerosis, autoimmune thyroiditis, bronchitis, cholecystitis,  
CC contact dermatitis, Crohn's disease, emphysema, erythroblastosis fetalis,  
CC glomerulonephritis, Good pasture's syndrome, gout, Grave's disease,  
CC Hashimoto's thyroiditis, multiple sclerosis, myasthenia gravis,  
CC osteoarthritis, osteoporosis, pancreatitis, psoriasis,  
CC Reiter's syndrome, arthritis, scleroderma, Sjogren's syndrome, systemic  
CC lupus erythematosus, ulcerative colitis, uveitis, Werner syndrome,  
CC complications of cancer, haemodialysis, and extracorporeal circulation,  
CC viral, bacterial, fungal parasitic, protozoal, and helminthic infections,  
CC trauma, or cancer. CME, or its catalytic or immunogenic fragment, is  
CC useful for drug screening.

XX Sequence 529 AA;

Query Match 68.7%; Score 1924.5; DB 21; Length 529;  
Best Local Similarity 69.9%; Pred. No. 1.5e-197;  
Matches 369; Conservative 47; Mismatches 109; Indels 3; Gaps 2;

```

OY 3 SDRALVFLLOLFC-VGCGFCGKVLVPCDMSHMLNKKVLEELIVGHEVTLTHSKP 61
DB 2 smktsaillilqscyscgkvlwptefshmmiklildygrhevtlaasas 61
OY 62 SLIDRRPALKFEVVMPCDRTENEFPVDALN-VLUGLSWOSVIRLNDFEVRIG 119
DB 62 isfipnstcltefvysltktefedilqlykrwaelpkdtwysfysqelmtwfind 121
OY 120 TLKMCSEFIYNOTLAKKLOETNYDWLIDVIPCGLMAELAVPVLTLRISVGNME 179
DB 122 ilkkfcdvsnkklmkkgserfdvdladavpfgellbellkkipvyltrfspyrate 181
OY 180 RSGGKLPAPLSYVPMPTGLDRTMLEKRNKMSLVLFHFWIODYDHFHEEFSKALG 239
DB 182 khsqgllfpssyvpvmseisdqmtiterkmllylylefqifomkkwdqfysevjg 241
OY 240 RPTTLCETVGKAEIWLIRTYWDEFPOPYQPNFEFVGLHCPRKAKLPKMEFVQSSGE 299
DB 242 rpttletnakadiwlirnywdfqfbrpdlipveifvgghckpapaklpkemeefvsgsg 301
OY 300 DGLVFSLSGLFONVTEKANIITASALAOIPQKVLWRKRGKRPSTLGANTRLYDWTPQND 359
DB 302 ngvvvslsgmvsntseetanviasalakipqkvlwrfqngkptlglitrllykvpqnd 361
OY 360 ILGHPRTKAFITGNGNGIYEATYHGVPMVGPVIFGDDOINDIAHMAKAAVETNKTYMT 419
DB 362 llyhpktkafitlghgmnglyealyngvpmvypifgqldidiahmakagaaveinlktnt 421
OY 420 SEDLALRLRTVTTSSYKENAMKLSRIHNDOPKPLDRAVVFVIEFVRRKGAKHLSAAH 479
DB 422 sedllralrtvtltdssykenamklsrihndgvpkdravvfietfmrkghkhltsaah 481
OY 480 DLTWFOHYSIDVIGLTLTVATAIFLFTGFLFSCQKFNKTKIEKRE 527
DB 482 dltwfhysidvlgfltlcvataiflftkclfcgckfnktrkiekre 529

```

RESULT 2  
ID AAM47126  
XX AAM47126 standard; Protein: 530 AA.

AC AAM47126;

XX 26-MAY-1998 (first entry)

DE Uridine diphospho-glucuronosyltransferase 2B17 (UGT2B17) enzyme.

KW Uridine diphospho-glucuronosyltransferase 2B17; UGT2B17; catalyse;  
XX androstereone; androstereone-glucuronic acid; androgen; enzyme.

XX Homo sapiens.

XX MO9744466-A1.

XX 27-NOV-1997.

XX 16-MAY-1997; 97WO-CA00328.

XX 17-MAY-1996; 96US-0649319.

XX (ENDO-) ENDORECHERCHE INC.

XX Beaulieu M, Belanger A, Hum DW, Levesque E;

XX WPI; 1998-018520/02.

XX N-PSDB: AAV15900.

PT DNA encoding uridine di:phospho:glucuronosyl:transferase 2B17 -

PT which catalyses conversion of androstereone to

XX androstereone-glucuronic acid

XX Claim 16; Pages 4-6; 53pp; English.

XX This is the enzyme uridine diphospho-glucuronosyltransferase 2B17  
CC (UGT2B17). This novel enzyme catalyses the conversion of androstereone  
CC to androstereone-glucuronic acid. The UGT2B17 can be used to detect  
CC anti-UGT2B17 antibodies. The antibody can be used to detect a localised  
CC concentration of UGT2B17 or an alteration in androgen activity. The  
CC UGT2B17 can also be used to alter the concentration of an androgenic  
CC compound in a tissue, specifically dihydrotestosterone. An isolated  
CC nucleotide sequence comprising at least 30 consecutive nucleotides from  
CC the coding region of the 2107 base pair sequence, or its complement can  
CC be used to block the synthesis of UGT2B17, e.g. an expression disrupting  
CC sense or antisense fragment, or as a probe for a UGT2B17 coding sequence.

XX Sequence 530 AA;

Query Match 61.8%; Score 1732; DB 19; Length 530;  
Best Local Similarity 61.5%; Pred. No. 7.1e-177;  
Matches 326; Conservative 74; Mismatches 112; Indels 18; Gaps 3;

```

OY 9 VFLLLOLFC-VGCGFCGKVLVPCDMSHMLNKKVLEELIVGHEVTLTHSKPSIDIR 67
DB 8 vflmlqscysfsgscgkvlwptefshmmiklildygrhevtlaasasallvns 67
OY 68 KRSALKFEVVMPCDRTENEFPVDALNVLPGLSWOSVIRLNDFEVRIGRTLK----- 123
DB 68 krsalkfevvmprtdrteneifvdlalnvlpglstwosvirlndefvrigrtlk----- 123
OY 124 -----MCSFIYNOTLAKKLOETNYDWLIDVIPCGLMAELAVPVLTLRISVGN 177
DB 121 sdynklcedavlnlkmkldeskdvdldavnpcegellaellnlpfllyslrfsvgt 180
OY 178 MERSCGKLPAPLSYVPMPTGLDRTMLEKRNKMSLVLFHFWIODYDHFHEEFSKALG 237
DB 181 vekngggflfpssyvpvmseisdqmtiterkmllylylefqifomkkwdqfysevjg 240
OY 238 LGRPTTLCETVGKAEIWLIRTYWDEFPOPYQPNFEFVGLHCPRKAKLPKMEFVQSSGE 297
DB 241 lgrpttletnakadiwlirnywdfqfbrpdlipveifvgghckpapaklpkemeefvsgsg 300
OY 298 GEDGIVFSLSGLFONVTEKANIITASALAOIPQKVLWRKRGKRPSTLGANTRLYDWTPQ 357
DB 298 gedgivfslsglfnvteekaniitasaalaoipokvlwrkrgkrrpstlgantrlwdtpq 357

```





```

Db 182 khsqgfifpssypvymseIdqmtfmervkmmlylvIdftwfefIdmqkwdqfyseIv 241
QY 240 RPTTLCTVGKAEIWLIRTWDEFPQYQPNFEVGGIHCCKPAKALPKEMENFVQSSGE 299
Db 242 rpttsetmgkadwllrnswnifqfypIlpnvdfvggIlnckpkpIpkemedfvsgsge 301
QY 300 DGIWVFISLGFQWVEKKNITIASALAOIPOKVLWRYKCKKPKETLGANTRLVYMTFOND 359
Db 302 ngvvfifsgmvsmtceeranvlasalaqIpoKvIwrfIdgnkpcIdIlntrIykwIpnq 361
QY 360 ILGHPKTKAEITHGNGMGIYEALYHGVPMGVPIFGDOLDNIAMKAKGAIVEINFKTM 419
Db 362 llghpktraftIhgnganglyealyhgIpmvgIpfIdagpnlIamkergaavrvdfnms 421
QY 420 SEDILRLRLVITDSSKEMAMLSRIHNDQPKRPLDRAVFIWFVNRHGAKHLRSAH 479
Db 422 stIdlnalkrvIdnpskyenvmkIsrIqhdpvIdravfIefvImthbgakhlrvaaH 481
QY 480 DLWFOHYSIDVIGFLTLVATAIFLFTKCFLEFSCOKENKTRK 522
Db 482 dlwfygshIdvIvgIflvcaIvIflvkcclIcfwktark 524

RESULT 6
ID AAR26153 standard; Protein: 533 AA.
AC AAR26153;
XX
XX 27-JAN-1993 (first entry)
XX
DE HUG-Br1.
XX
XX Billirubin; UDP-glucuronosyltransferase; HUGBr1; HUGBr2;
KW monoglucuronide; diglucuronide.
XX
OS Homo sapiens.
XX
FH key
FT Region 10..20 location/Qualifiers
FT /note= "putative membrane-insertion signal"
FT Region 491..507
FT /note= "putative membrane-anchoring peptide"
FT Modified-site 102
FT /note= "predicted Asn-linked glycosylation site"
FT Modified-site 295
FT /note= "predicted Asn-linked glycosylation site"
FT Modified-site 347
FT /note= "predicted Asn-linked glycosylation site"
FT MISC-difference 158
FT /note= "feature not labelled in specification"
FT MISC-difference 181
FT /note= "feature not labelled in specification"
FT MISC-difference 228
FT /note= "feature not labelled in specification"

WO9212987-A.
XX
PD 06-AUG-1992.
XX
PF 10-JAN-1992. 92MO-US00282.
XX
XX 10-JAN-1991. 91US-0639453.
XX
PA (USSH ) US DEPT HEALTH & HUMAN SERVICE.
XX
PI Owens IS, Ritter JK;
XX
XX MPI: 1992-284593/34.
XX
XX N-PSDB; AA027369.
XX
XX
XX Isolated gene locus UGT1. DNA segments and diagnostic probes -
PT for diagnosing Gilbert's disease and Crigler-Najjar syndrome

```

```

PT types I and II
XX
XX PS Disclosure; Fig 9A-I; 99pp; English.
XX
XX Two human liver bilirubin UDP-glucuronosyltransferase cDNAs have
CC been isolated. They are referred to as HUGBr1 (AA027369) and HUGBr2
CC (AA027370) (Ritter, et al., J. Biol. Chem. 266:1043-1047 (1991)) and,
CC upon expression individually in COS-1 cells, encode isoforms that
CC catalyse the formation of the two bilirubin monoglucuronides and
CC the diglucuronide.
CC The cDNAs contain identical 3' ends (1469 bp in length) to each
CC other and to that of the human phenol transferase cDNA, HUGP1
CC (Harding et al., Proc. Natl. Aca. Sci. USA 85:8281 (1988)).
CC In contrast, they have unique 5' ends.
XX
SQ Sequence 533 AA;

Query Match 40.0%; Score 1120; DB 13; Length 533;
Best Local Similarity 45.0%; Pred. No. 3.8e-111;
Matches 232; Conservative 88; Mismatches 186; Indels 10; Gaps 6;

QY 15 LFCV---GCGFCGKVLVPCDMSHMLNVKYLEELVRGHEVVLTHSKPSLDYRKPRA 71
Db 16 llcvlgpvnshagklllpvdgshwsmIgaIqIgrgheIvLa---pdaIyIrdga 72
QY 72 L-KFEVVMPODRTENEIIFDALNVLPGLISTWQSVYIKNDFEVEIRGLKMKCESFIY 130
Db 73 fytIktyrpfqredvkesfvsIghnvfendsIqrvIkkykIkKkdsamIsgcsIhIh 132
QY 131 NQTLMKRLOETNYVMILIDVYIPCGDLMAELANPFLTLRISVGMMERSCGKLPAPLS 190
Db 133 nkelmaslaessIdvmltdpIpcspIvaqIIsrIpfvIln-aiPcsIfeateqcpnfS 191
QY 191 YVPVMTGLIDRMFTLEVRKNSMLSVFHFWDYDVFHFWEFEFSKALGRPTTLCTETVGK 250
Db 192 yvrpIshsdhmtfIqrvkmIlaIafsqnf-IcdvvsyptIaIaseIqrevtvqdIIs 250
QY 251 AEIWLIRTWDEFPQYQPNFEVGGIHCCKPAKALPKEMENFVQSSGEQIVVFSISGL 310
Db 251 asvwlfsrdIvkdypIpmnvfvvgIlnclhnpIsqefeaYInaIsehgIvIvIsIesm 310
QY 311 FQNTTEKANITIASALAOIPOKVLWRYKCKKPKETLGANTRLVYMTFONDILGHPKTKAFI 370
Db 311 vsetIpekkaIadaIqkIpoKvIwrfIgcIpsIlnanclIvkwIpnqdlIghpmtIaIh 370
QY 371 THGGMNGIYEALYHGVPMGVPIFGDOLDNIAMKAKGAIVEINFKTMSEDLRALRTV 430
Db 371 thgshgvyesIcngvpmvmpIlgdqmdnakmetkgagvtIlnvIamtedIenaqkav 430
QY 431 ITDSSYKEMAMLSRIHNDQPKRPLDRAVFIWFVNRHGAKHLRSAHDLTWFOHYSID 490
Db 431 IndksyenImrIessIhIkdIrvpeIdIavfweIvmthbgakhlrpaahdltwyqyIsId 490
QY 491 VIGFILTCAVTAIFLFTKCFLEFSCOK-FNKTRKIEK 525
Db 491 vIgfIlIavIltvaIftfkcaayrIkclqIkkgyrvKk 526

RESULT 7
ID AAR26154 standard; Protein: 534 AA.
XX
XX AAR26154;
XX
XX 27-JAN-1993 (first entry)
XX
XX HUG-Br2.
XX
XX
XX Billirubin; UDP-glucuronosyltransferase; HUGBr1; HUGBr2;
KW monoglucuronide; diglucuronide.
XX
XX Homo sapiens.

```

```

XX Key Location/Qualifiers
FH Region 12..22
FT /note="putative membrane-insertion signal"
FT Region 492..508
FT /note="putative membrane-anchoring peptide"
FT Modified-site 348
FT /note="predicted Asn-linked glycosylation site"
FT Misc-difference 282..285
FT /note="residues encoded by TGCCAACGGAGC!"
XX WO9212987-A.
PD 06-AUG-1992.
XX 10-JAN-1992; 92WO-US00282.
XX 10-JAN-1991; 91US-0639453.
XX (USSH ) US DEPT HEALTH & HUMAN SERVICE.
XX Owens IS, Rilter JK;
XX WPI: 1992-284593/34.
XX N-PSDB; AAQ27369.
XX Isolated gene locus UGT1, DNA segments and diagnostic probes -
XX for diagnosing Gilbert's disease and Crigler-Najjar syndrome
XX types I and II
XX Disclosure; Fig 9A-I; 99pp; English.
XX Two human liver bilirubin UDP-glucuronosyltransferase cDNAs have
XX been isolated. They are referred to as HUGB1 (AAQ27369) and HUGB2
XX (AAQ27370) (Rilter, et al., J. Biol. Chem. 266:1043-1047 (1991)) and,
XX upon expression individually in COS-1 cells, encode isoforms that
XX catalyse the formation of the two bilirubin monoglucuronides and
XX the diglucuronide.
XX The cDNAs contain identical 3' ends (1469 bp in length) to each
XX other and to that of the human phenol transferase cDNA, HUGP1
XX (Harding et al., Proc. Natl. Aca. Sci. USA 85:8281 (1988)).
XX In contrast, they have unique 5' ends.
XX Sequence 534 AA:
SQ
Query Match 37.8%; Score 1058.5; DB 13; Length 534;
Best Local Similarity 41.9%; Pred. No. 1.5e-104;
Matches 227; Conservative 86; Mismatches 172; Indels 57; Gaps 8;
QY 11 LLLQLFCVGGCGFKVYVWPCDMSHMLNVKVLLEELIVRGHEVTVLT-----HKKP--- 61
DB 16 LLLLlLVqpaesgkvlvprtdgspwlsmealrelharghgvavltprgeyahgeekff 75
QY 62 SLIDYKPSALK-----FEVYHMPQDRTEENEFVDLALNVLPGISTQSVI 108
DB 76 cltayavpwtqkefdvltlygtgffetehtllkrysrmaimnvsI----- 122
QY 109 KLNDFFVEIRGTLKMKCESFIVNOTLKKLQENINVMALIDYTPCGDLMAELIAPVYL 168
DB 123 -----ahrcvaelllmeallrhlnatstidvltcpnlgavtlakylsipaaf 171
QY 169 TURISVGMMERSCGRLPAPLSTVPVPMGTLDRMFELRRVKNM-----LSVLFHWID 224
DB 172 fwrty-ipcldlfdqtcgpcpsysyiplklitndmftfgrvknmyplalsyichfsap 230
QY 225 YVHWEERYYSALGRPTLCEVGAELIWLIRTYWDEFFPOPOPNFEFGVGLHCKPAK 284
DB 231 Y-----aslaselfgrevlvqdlssaswvlfrsdftvkdpimpnmvffigincangk 285
QY 285 ALPKEMENFVQSSGEGIVVFSIGSLFQWVPEKANIITASALAOIPQKYLWKYKGGKST 344
DB 286 pslgqfeayinasgehgvlvftslsmvselpekkamataladlqktpvtlvrtlygtlrpsn 345

```

```

QY 345 LGANRLYDWIPONDLLGPKTKAPITLGGMNGIYEATIGHVPMYGVPIFGDLNIAHM 404
DB 346 lanncllkwlpqndllghpmtralfitlsgnsyngvscngvpmvmpplfgqmdnakrm 405
QY 405 KAKGAAVEINERTWTSDDLRLALRTVITDSSYKENAMRLSRINHDOPKPLDRAVFWIEF 464
DB 406 elkgagvltlnvlemtsedlmaeqkavindksykenimrlsslhkdrprepidlavfwef 465
QY 465 VNRHKGAKHLRSAADLTLWPFQHSYSDVIGELLITCATALFELFKCFLESCOK-FKTRKI 523
DB 466 vmrhkgaphltpaandltwygshdvlvllavlltvalitfkccagyrkclgkgrv 525
QY 524 EK 525
DB 526 KK 527
RESULT 8
AA57100
ID AA57100 standard; Protein: 245 AA.
XX
AC AA57100;
DT 28-FEB-2000 (first entry)
XX
DE UDP-glucuronosyltransferase 1 (UGT1) exons 2-5 amino acid sequence.
XX
KW uridine diphosphate-glucuronosyltransferase 1; UGT1; polymorphism; probe;
KW glucuronic acid; Crigler-Najjar syndrome; Gilbert syndrome; jaundice;
KW unconjugated hyperbilirubinaemia; drug metabolism; transgenic animal;
KW pharmacogenetic screening; diagnose.
XX
OS Homo sapiens.
XX
PN WC0957322-A2.
PD 11-NOV-1999.
PF .04-MAY-1999; 99WO-US09702.
XX
PR 07-MAY-1998; 98US-0084807.
XX
PA (AXYS-) AXYS PHARM INC.
XX
PI Penny L, Galvin M;
XX
DR WPI: 2000-052981/04.
DR N-PSDB: AA245118.
XX
PT New nucleic acid representing polymorphisms in the human uridine
PT diphosphate glucuronosyltransferase gene, used for diagnosis and
PT evaluation of drug metabolism
XX
PS Examples; Page 44-45; 63pp; English.
XX
CC AA57092-57100 are the amino acid sequences of exons 1A-1J of human
CC uridine diphosphate-glucuronosyltransferase 1 (UGT1). The UGTs are a
CC family of enzymes that catalyse the glucuronic acid conjugation of a
CC wide range of endogenous and exogenous substrates including phenols,
CC alcohols, amines and fatty acids. Many of the reactions catalysed by
CC UGTs result in toxic substances being converted to compounds which are
CC more water soluble and are excreted. The invention relates to and
CC identifies UGT1 polymorphisms (AA245004-245041). The polymorphism
CC sequences are useful as probes for detecting UGT1 locus polymorphisms,
CC indicative of altered UGT1 expression or activity. These polymorphisms
CC are associated with Crigler-Najjar and Gilbert syndromes (unconjugated
CC hyperbilirubinaemia) and drug metabolism. The genotyping of the UGT1 gene
CC is used to predict the rate of metabolism of UGT1 substrates, possible
CC drug-drug interactions and adverse side effects (i.e. to optimize drug
CC dosage), and to screen for diseases caused by exposure to toxins and to
CC study the effects of polymorphisms on enzymatic activity. The UGT1
CC sequences, including polymorphisms, can also be used to produce the

```

CC corresponding protein (or its fragments) or to generate transgenic  
CC animals or modified cells e.g. for pharmacogenetic screening.

50 Sequence 245 AA;

Query Match	27.0%;	Score 755.5;	DB 21;	Length 245;
Best Local Similarity	59.7%;	Pred. No. 1.5e-72;		
Matches 142;	Conservative 34;	Mismatches 61;	Indels 1;	Gaps 1;

QY 289 EMEFVQSSGEGDGIWFSLSGLFQNTTEEKANITIASALAOIPQKVLWRYKKKKPSTIGAN 348  
| : : ||| ||||| : : || ||| : ||| |||| : |||  
Db 1 efaaylnasgehgivfslsgsmvseipekkamaladalgktpqvtlwyrgttrpsnlann 60

QY 349 TRLDWIPQNDLGRPKAFITHGSGNGIYEALYHGVMGVPRFGDQDLNIAHHKAKG 408  
| | : | | | | | : | | | | : | | | | : | | : | |  
Db 61 TLVkwlpqndllgpmtratlthagshgyeslcnvgymvmmprlfgdqmdnakrmetkq 120

Qy 409 AANEINFKMTSEDLRALRIVYITDSSYKENARLSTRHHDPKPLDRATVEWIEFMRH 468  
| : | ||||| | : | | ||||| |||| : | : ||||| |||| : |||||  
Db 121 agvlinvlemtsedlenalkavindksyenimrisslhkdrpepldlavfwefmrh 180

QY 469 KGAKHLRSAHDLTWFGHSIDVIGLTLTCAATAIFLFKCFLEFSÖK-FNKTRKIEK 525  
||| ||| |||||||:::||| | : ||| : || ::|  
Db 181 kgaphlrpaandltwygysldvlgfllavlltvaftfcckcaygyrkclgkkgvrkk 238

RESULT	9
AAB24025	
ID	AAB24025 standard; Protein; 523 AA.

AC AAB24025;

DT 25-JAN-2001 (first entry)

DE Human PRO1780 protein sequence SEQ ID NO:13.

KM Human; tumour; diagnosis; neoplastic disease; proliferation; cancer; identification; tumourigenesis; anticancer; detection.

OS Homo sapiens.

PN WO200053750-A1.

PD 14-SEP-2000.

PF 02-DEC-1999; 99WO-US28551.

PR 08-MAR-1999; 99WO-US05028.

PR 29-OCT-1999; 99US-0162506.

PR 01-DEC-1999; 99WO-US28634.

PA (GETH ) GENENTECH INC.

PI Botstein D, Goddard A, Gurney AL, Roy MA, Watanabe CK, Wood WI;

DR WPI; 2000-594320/56.

[illegible]

PT the growth of tumors in mammals, and to identify inhibitors of PRO

XX  
XX  
1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840[illegible]

CC protein (I) selected from: PRO381, PRO1269, PRO1410, PRO1755, PRO1780,  
CC PRO3434, PRO1927, PRO3567, PRO1295, PRO1293, PRO1303, PRO3444, PRO4354, CC

CC anticancer activity and can be used to diagnose tumors in mammals, by  
CC detecting complex formation when the antibody is contacted with test  
CC

CC cells. Increased expression of genes encoding (I) can also be detected  
CC to diagnose tumours. Agents which inhibit the activity of (I),  
CC especially the antibodies, or an antisense oligonucleotide which,  
CC hybridises to genes encoding (I), can be used to inhibit tumour growth,  
CC preferably by inducing cell death. Methods from the present invention  
CC can be used to identify compounds which inhibit the biological activity  
CC of (I). AAC58019 to AAC58102 represent PCR primers and hybridisation  
CC probes used in examples from the present invention for human PRO  
CC sequences. AAC58103 to AAC58122 and AAB24021 to AAB24040 represent human  
CC PRO polynucleotide and protein sequences given in the exemplification of  
CC the present invention.

50 Sequence 523 AA;

Query Match	25.58;	Score 714.5;	DB 21;	Length 523;
Best Local Similarity	33.98;	Pred. No. 1.3e-67;		
Matches 172;	Conservative 93;	Mismatches 208;	Indels 35;	Gaps 13;

```

Oy      34 SHMLNKKVILEELIVRGHEVYLLTHSK-PSLIDYRKPSALKFEVHH-MPDDRTENEIEIF 90
      ||::: : : ||||: : : ::| : : | :
Db      34 shyllmdrvsllqdhghnvtlnhkrgrpfmddkkek-syqvtswlapedhqrctkks 92

```

```

OY      | | | : : : | | | : | : : :
91 VDIAL-NVEGLSTWOSVIKLNDFEVEIRGTLKMKCESFYIQTLMKKLOETINYDWALID 1493
Db      | | | : : : | | | : | : : :
93 fdfleetlgrgkfenlnvle-----laqcshtfnrkdlmslknfenfawive 1455

```

```

QY      150  PVIPEGDMLMELLAVPEVLTIRISVGGMNRSCKLPALPSYVPVPMGTLDRTMFLERV 209
          | : | | | | | | | | | | | | | | | | | | | | | | | | | |
Db      146  tfdycpfllekkgkprivalstsf--gslef---lpipisypvprfsltdhmdfwgrv 201

```

```

QY      210 KNSM$VL$----HFWDIDYD$HFMEE$YSKALGRPTTLCETVGAEIWLIRTYWDEEP    265
          || :: | :| | | | | :|||:| : : |:|
Db      202 knfmffsfcrqgmngstfahltikenhfe---gsrplvshlllkaelwifnsdfadfa    258

```

259 rpllpntvvvglmekpikpypqdlenfiafkfsgsfvltlgsnmvntcqnpeflkemm 318

```

321 11ASALADLEPRVLMWKIR--"GAKPSISGIANIKIRUMLEPNULDSHPKINAFIIRDSMG 374
      |::||:::| |::|::|::|::|::|::|::|::|::|::|::|::|::|::|::|::|
Db 319 ----afahlpqgviwkcqscshpdkvhaanavkivdwlpsdllaipssirllfvthgqns 374

```

Db 375 imeatqhgvpmyiprlfgdqpemvrveakkfysysiqllkklkaetalkmqimedkryx 434

Db 435 saavaasvllrshplptqrlvgwidhnlqtgcatlhkpyvfqgpwhnegylfdvfvfllg 494

495 1t1gt1wlcgk1lgmavwmlrgarkvke 522

RESULT	10
AAV99419	
ID	AAV99419 standard; Protein; 523 AA

AC	AAV99419;
XX	
DT	08-AUG-2000 (first entry)

DE	Human PRO1780 (UNQ842) amino acid sequence SEQ ID NO:282
XX	
KW	Human: PRO polypeptide; membrane bound protein; receptor

XX  
OS  
XX  
PN

Homo sapiens.

WC0000132708-A2

PD 09-MAR-2000.  
 XX 01-SEP-1999; 99WO-US20111.  
 XX  
 PR 01-SEP-1998; 98US-0098716.  
 PR 01-SEP-1998; 98US-0098749.  
 PR 01-SEP-1998; 98US-0098750.  
 PR 02-SEP-1998; 98US-0098803.  
 PR 02-SEP-1998; 98US-0098821.  
 PR 02-SEP-1998; 98US-0098843.  
 PR 09-SEP-1998; 98US-0099536.  
 PR 09-SEP-1998; 98US-0099596.  
 PR 09-SEP-1998; 98US-0099598.  
 PR 09-SEP-1998; 98US-0099602.  
 PR 10-SEP-1998; 98US-0099642.  
 PR 10-SEP-1998; 98US-0099741.  
 PR 10-SEP-1998; 98US-0099754.  
 PR 10-SEP-1998; 98US-0099763.  
 PR 10-SEP-1998; 98US-0099792.  
 PR 10-SEP-1998; 98US-0099808.  
 PR 10-SEP-1998; 98US-0099812.  
 PR 10-SEP-1998; 98US-0099815.  
 PR 15-SEP-1998; 98US-0100385.  
 PR 15-SEP-1998; 98US-0100388.  
 PR 15-SEP-1998; 98US-0100390.  
 PR 16-SEP-1998; 98US-0100584.  
 PR 16-SEP-1998; 98US-0100627.  
 PR 16-SEP-1998; 98US-0100661.  
 PR 16-SEP-1998; 98US-0100662.  
 PR 16-SEP-1998; 98US-0100664.  
 PR 17-SEP-1998; 98US-0100683.  
 PR 17-SEP-1998; 98US-0100684.  
 PR 17-SEP-1998; 98US-0100710.  
 PR 17-SEP-1998; 98US-0100711.  
 PR 17-SEP-1998; 98US-0100919.  
 PR 18-SEP-1998; 98US-0100930.  
 PR 18-SEP-1998; 98US-0100848.  
 PR 18-SEP-1998; 98US-0101014.  
 PR 18-SEP-1998; 98US-0101014.  
 PR 18-SEP-1998; 98US-0101068.  
 PR 18-SEP-1998; 98US-0101068.  
 PR 22-SEP-1998; 98US-0101279.  
 PR 23-SEP-1998; 98US-0101279.  
 PR 23-SEP-1998; 98US-0101471.  
 PR 23-SEP-1998; 98US-0101472.  
 PR 23-SEP-1998; 98US-0101474.  
 PR 23-SEP-1998; 98US-0101475.  
 PR 23-SEP-1998; 98US-0101476.  
 PR 23-SEP-1998; 98US-0101477.  
 PR 23-SEP-1998; 98US-0101479.  
 PR 24-SEP-1998; 98US-0101738.  
 PR 24-SEP-1998; 98US-0101741.  
 PR 24-SEP-1998; 98US-0101743.  
 PR 24-SEP-1998; 98US-0101915.  
 PR 24-SEP-1998; 98US-0101916.  
 PR 29-SEP-1998; 98US-0102207.  
 PR 29-SEP-1998; 98US-0102240.  
 PR 29-SEP-1998; 98US-0102307.  
 PR 29-SEP-1998; 98US-0102307.  
 PR 30-SEP-1998; 98US-0102331.  
 PR 30-SEP-1998; 98US-0102484.  
 PR 30-SEP-1998; 98US-0102487.  
 PR 30-SEP-1998; 98US-0102570.  
 PR 30-SEP-1998; 98US-0102571.  
 PR 01-OCT-1998; 98US-0102684.  
 PR 01-OCT-1998; 98US-0102687.  
 PR 02-OCT-1998; 98US-0102965.  
 PR 06-OCT-1998; 98US-0103258.  
 PR 06-OCT-1998; 98US-0103449.  
 PR 07-OCT-1998; 98US-0103314.  
 PR 07-OCT-1998; 98US-0103315.  
 PR 07-OCT-1998; 98US-0103328.  
 PR 07-OCT-1998; 98US-0103395.

PR 07-OCT-1998; 98US-0103396.  
 PR 07-OCT-1998; 98US-0103401.  
 PR 08-OCT-1998; 98US-0103633.  
 PR 08-OCT-1998; 98US-0103678.  
 PR 08-OCT-1998; 98US-0103679.  
 PR 14-OCT-1998; 98US-0103711.  
 PR 20-OCT-1998; 98US-0104257.  
 PR 20-OCT-1998; 98US-0104987.  
 PR 20-OCT-1998; 98US-0105000.  
 PR 21-OCT-1998; 98US-0105002.  
 PR 22-OCT-1998; 98US-0105104.  
 PR 22-OCT-1998; 98US-0105169.  
 PR 26-OCT-1998; 98US-0105266.  
 PR 26-OCT-1998; 98US-0105693.  
 PR 26-OCT-1998; 98US-0105694.  
 PR 27-OCT-1998; 98US-0105807.  
 PR 27-OCT-1998; 98US-0105881.  
 PR 27-OCT-1998; 98US-0105882.  
 PR 27-OCT-1998; 98US-0106062.  
 PR 28-OCT-1998; 98US-0106062.  
 PR 28-OCT-1998; 98US-0106029.  
 PR 28-OCT-1998; 98US-0106030.  
 PR 28-OCT-1998; 98US-0106033.  
 PR 28-OCT-1998; 98US-0106033.  
 PR 28-OCT-1998; 98US-0106178.  
 PR 29-OCT-1998; 98US-0106248.  
 PR 29-OCT-1998; 98US-0106384.  
 PR 29-OCT-1998; 98US-0108500.  
 PR 30-OCT-1998; 98US-0106464.  
 PR 03-NOV-1998; 98US-0106856.  
 PR 03-NOV-1998; 98US-0106902.  
 PR 03-NOV-1998; 98US-0106905.  
 PR 03-NOV-1998; 98US-0106919.  
 PR 03-NOV-1998; 98US-0106932.  
 PR 10-NOV-1998; 98US-0106934.  
 PR 10-NOV-1998; 98US-0107783.  
 PR 17-NOV-1998; 98US-0108775.  
 PR 17-NOV-1998; 98US-0108779.  
 PR 17-NOV-1998; 98US-0108787.  
 PR 17-NOV-1998; 98US-0108788.  
 PR 17-NOV-1998; 98US-0108801.  
 PR 17-NOV-1998; 98US-0108802.  
 PR 17-NOV-1998; 98US-0108806.  
 PR 17-NOV-1998; 98US-0108807.  
 PR 17-NOV-1998; 98US-0108825.  
 PR 18-NOV-1998; 98US-0108848.  
 PR 18-NOV-1998; 98US-0108849.  
 PR 18-NOV-1998; 98US-0108850.  
 PR 18-NOV-1998; 98US-0108851.  
 PR 18-NOV-1998; 98US-0108852.  
 PR 18-NOV-1998; 98US-0108858.  
 PR 18-NOV-1998; 98US-0108904.

(GETH ) GENENTECH INC.

Baker K, Goddard A, Gurney AL, Smith V, Watanabe CK, Wood WI;  
 WPI: 2000-237871/20.  
 DR N-PSDB; AAA37101.

XX  
 PT New mammalian DNA sequences encoding transmembrane, receptor or  
 PT secreted PRO polypeptides, useful for screening of potential peptide or  
 PT small molecule inhibitors of the relevant receptor/ligand interactions  
 XX  
 PS Claim 12; Fig 160; 773pp; English.

CC AAA37022 to AAA37144 encode the new isolated human transmembrane,  
 CC receptor or secreted PRO polypeptides given in AA99340 to AA99462. The  
 CC transmembrane and receptor PRO proteins can be used for screening of  
 CC potential peptide or small molecule inhibitors of the relevant  
 CC receptor/ligand interactions. The polypeptides and nucleotide sequences  
 CC encoding then have various industrial applications, including uses as



CC pharmaceutical and diagnostic agents, AAA37145 to AAA37330 represent  
CC PCR primers and hybridisation probes used in the isolation of the PRO  
CC polypeptides from the present invention.

XX Sequence 523 AA;

Query Match 25.5%; Score 714.5; DB 21; Length 523;

Best Local Similarity 33.9%; Pred. No. 1.3e-67;

Matches 172; Conservative 93; Mismatches 208; Indels 35; Gaps 13;

34 SHMLNKKVILEELVIRGHEVTVLTHSK-PSLIDYKPKALFEVYH--MPQDRENEEIF 90  
Db 34 shylmdtvsqllqdhgvnvtlnkhkrgpmpdfkkek-syqvswlapedhqrfeiks 92  
Qy 91 VDIAL-NVLPGLSTWOSYIKINDFEVEIRGTLKMMCESFIYNOTLMKKLOETNYDVMLID 149  
Db 93 fdfleeetlgrgkfeenlnvley-----lajqcsfhnrdkmdsknefmvive 145  
Qy 150 PVIRCGDLMAELLAVPVLTIRISYGVGMERSCKLPAPLSVVPVMTGLTDRMTLEERV 209  
Db 146 tfdycpfliaeklgkpfvaiststf-gsliefg---lpiplyvpyfrslltchmdfwgrv 201  
Qy 210 KNSMLSVLF---HFVIDDYHMEEFYSKALGPTTLCTGVGAELTWRITVWDEFFP 265  
Db 202 knlmlfscrrqdmgstfndcltkehfe---gsrpylshlllkaelwflnsdafa 258  
Qy 266 QPYOPNFEFVGGLHCKPAKALPKEMENFVOSGEGDIYVFSLSLF---QN--VTEERKAN 320  
Db 259 rpllpntvvgvlgmekrkpkpvgdlenfakfgdsqfvlvlgsvntcqppeitkemm 318  
Qy 321 IIASALAOIPQKVLIRYK---GKKPSTLGANTRLYDWIPQNDLGHPRTKAFTTHGMNG 377  
Db 319 ---afahlpqgvitwkcqshwpkdvhaanvkivdwipgsdillahpsirllfvthgqns 374  
Qy 378 IYEATYHGVPMVGPVIFGDODNTAHMKAGAAVEINKTMSSEDLALAKRVITDSSYK 437  
Db 375 imeaighvpmvgyiplfdgqpenmryeaakfgyvsigklkkaelalkmqimedkryk 434  
Qy 438 ENAMRLSRINHDQVPKPLDRAVFIETPMRRHKGAKHLRSAHADLTWFOHYSIDVIGFLIT 497  
Db 435 saavaasviltshpsrptqrlvgwidhvlqtcgathlkpryvfqrpwheqylfdvfvllg 494  
Qy 498 CVATAIFLFTKCFLESCQKFNKTRIEK 525  
Db 495 ltlgtlwlgckllgmavwvlrgarkvke 522

# RESULT 11

AA66168 standard; protein; 523 AA.

AC AAB66168;

DT 02-APR-2001 (first entry)

DE Protein of the invention #80.

KW Secreted; transmembrane; gene therapy.

OS Unidentified.

XX W0200078961-A1.

XX 28-DEC-2000.

PE 18-FEB-2000; 2000MO-US04342.

PR 23-JUN-1999; 99US-0141037.

PR 20-JUL-1999; 99US-0144758.

PR 26-JUL-1999; 99US-0145698.

PR 01-SEP-1999; 99MO-US20111.

PR 29-OCT-1999; 99US-0162506.

PR 30-NOV-1999; 99MO-US28313.  
PR 02-DEC-1999; 99MO-US28551.  
PR 16-DEC-1999; 99MO-US30095.  
PR 05-JAN-2000; 2000MO-US00219.  
PR 06-JAN-2000; 2000MO-US00376.

XX (GETH ) GENENTECH INC.

XX Baker KP, Botstein D, Desnoyers L, Eaton DL, Ferrara N, Fong S;  
XX Gao W, Goddard A, Godowski PJ, Grimaldi CJ, Gurney AL, Hillan KJ;  
XX Pan J, Paon NF, Roy MA, Smith V, Stewart TA, Tamas D;  
XX Watanabe CK, Williams PM, Wood WI;  
XX WPI; 2001-071395/08.

Secreted and transmembrane proteins and nucleic acids designated PRO,  
PT useful as hybridization probes, in chromosome and gene mapping and gene  
therapy -  
Claim 1; Fig 160; 787pp; English.

The present invention relates to secreted and transmembrane proteins.  
These proteins and the DNA encoding them may be used as hybridization  
probes, in chromosome and gene mapping and in the generation of  
anti-sense RNA and DNA. They may also be used to generate either  
transgenic animals or knockout animals which are in turn useful for  
development and screening of therapeutically useful reagents.  
The nucleic acids may also be used in gene therapy.

XX Sequence 523 AA;

Query Match 25.5%; Score 714.5; DB 22; Length 523;

Best Local Similarity 33.9%; Pred. No. 1.3e-67;

Matches 172; Conservative 93; Mismatches 208; Indels 35; Gaps 13;

34 SHMLNKKVILEELVIRGHEVTVLTHSK-PSLIDYKPKALFEVYH--MPQDRENEEIF 90  
Db 34 shylmdtvsqllqdhgvnvtlnkhkrgpmpdfkkek-syqvswlapedhqrfeiks 92  
Qy 91 VDIAL-NVLPGLSTWOSYIKINDFEVEIRGTLKMMCESFIYNOTLMKKLOETNYDVMLID 149  
Db 93 fdfleeetlgrgkfeenlnvley-----lajqcsfhnrdkmdsknefmvive 145  
Qy 150 PVIRCGDLMAELLAVPVLTIRISYGVGMERSCKLPAPLSVVPVMTGLTDRMTLEERV 209  
Db 146 tfdycpfliaeklgkpfvaiststf-gsliefg---lpiplyvpyfrslltchmdfwgrv 201  
Qy 210 KNSMLSVLF---HFVIDDYHMEEFYSKALGPTTLCTGVGAELTWRITVWDEFFP 265  
Db 202 knlmlfscrrqdmgstfndcltkehfe---gsrpylshlllkaelwflnsdafa 258  
Qy 266 QPYOPNFEFVGGLHCKPAKALPKEMENFVOSGEGDIYVFSLSLF---QN--VTEERKAN 320  
Db 259 rpllpntvvgvlgmekrkpkpvgdlenfakfgdsqfvlvlgsvntcqppeitkemm 318  
Qy 321 IIASALAOIPQKVLIRYK---GKKPSTLGANTRLYDWIPQNDLGHPRTKAFTTHGMNG 377  
Db 319 ---afahlpqgvitwkcqshwpkdvhaanvkivdwipgsdillahpsirllfvthgqns 374  
Qy 378 IYEATYHGVPMVGPVIFGDODNTAHMKAGAAVEINKTMSSEDLALAKRVITDSSYK 437  
Db 375 imeaighvpmvgyiplfdgqpenmryeaakfgyvsigklkkaelalkmqimedkryk 434  
Qy 438 ENAMRLSRINHDQVPKPLDRAVFIETPMRRHKGAKHLRSAHADLTWFOHYSIDVIGFLIT 497  
Db 435 saavaasviltshpsrptqrlvgwidhvlqtcgathlkpryvfqrpwheqylfdvfvllg 494  
Qy 498 CVATAIFLFTKCFLESCQKFNKTRIEK 525  
Db 495 ltlgtlwlgckllgmavwvlrgarkvke 522



CC untranslated region (UTR) of the mRNA because they are often obtained  
CC from oligo-dT primed cDNA libraries. Such ESTs are not well suited for  
CC isolating cDNA sequences derived from the 5' ends of mRNAs and even in  
CC those cases where longer cDNA sequences have been obtained, the full 5'  
CC UTR is rarely included. 5' ESTs are derived from mRNAs with intact 5'  
CC ends and can therefore be used to obtain full length cDNAs and genomic  
CC DNAs. 5' ESTs are also used in diagnostic, forensic, gene therapy and  
CC chromosome mapping procedures. They are used to obtain upstream  
CC regulatory sequences and to design expression and secretion vectors.

XX Sequence 78 AA:

Query Match 14.5%; Score 405; DB 21; Length 78;  
Best Local Similarity 100.0%; Pred. No. 1.1e-35;  
Matches 78; Conservative 0; Mismatches 0; Indels 0; Gaps 0;

OY 290 MEHFVSSGPDGIVFVSLGSLPQNVTEERKNIITASALAOIPQKVLNRYKGRKSTLGANT 349  
|||||  
Db 1 menfvgssgedgvlvslgslqnvteekaniiasalaqipkvlywrykgrkstlgant 60

OY 350 RLYDMIPQNDLGHPRKTK 367  
|||||  
Db 61 rlydwlipqndllghprtk 78

#### RESULT 14

AA29525  
ID AAY29525 standard; Protein; 129 AA.

XX AAY29525;

XX 13-OCT-1999 (first entry)

XX Human lung tumour protein Lr86-5 predicted amino acid sequence.

XX Human; lung tumour protein; therapy; diagnosis; lung cancer; vaccine;  
XX immunotherapy; detection; inhibition.

XX Homo sapiens.

XX WO938973-A2.

XX 05-AUG-1999.

XX 26-JAN-1999; 99MO-US01642.

XX 22-DEC-1998; 98US-0219245.

XX 28-JAN-1998; 98US-0015022.

XX 28-JAN-1998; 98US-0015029.

XX 18-MAR-1998; 98US-0040828.

XX 18-MAR-1998; 98US-0040831.

XX 23-JUL-1998; 98US-0122191.

XX 23-JUL-1998; 98US-0122192.

XX (CORI-) CORIXA CORP.

XX Erudakis TN, Lodes MJ, Mohamath R, Reed SG;

XX WPI; 1999-479187/40.

XX N-PSDB; AA207208.

XX Lung tumour specific polynucleotides for inhibiting the development  
XX of lung cancer

XX Example 2; Page 73; 171pp; English.

XX The present invention describes lung tumour specific polynucleotides  
XX and tumour antigens. AA207144 to AA207246 and AA208301 to AA208325  
XX represent specifically claimed polynucleotides, and AAY29486 to AAY29571  
XX represent amino acid sequences from the present invention. The lung  
XX tumour specific polynucleotides and polypeptides can be used in  
XX pharmaceutical compositions and vaccines to inhibit the development of

CC lung cancer. They can also be used to detect lung cancer in a patient.  
CC Probes and antibodies derived from the lung tumour sequences are useful  
CC in detection of lung cancer.

XX Sequence 129 AA:

Query Match 14.0%; Score 391; DB 20; Length 129;  
Best Local Similarity 58.1%; Pred. No. 7.6e-34;  
Matches 75; Conservative 18; Mismatches 36; Indels 0; Gaps 0;

OY 301 GIVFSLGSLPQNVTEERKNIITASALAOIPQKVLNRYKGRKSTLGANTRLVDMIPQNDL 360  
|||||  
Db 1 givfslgsmvseipekkavaiadalgktpgtvlywrytgrpsnlanntilvqwlipqndl 60

OY 361 LGHPKTKARITHCNMGIYEAITYHGVPMVGPPIFGPOLNINAHMKAKAIVEINFRTMTS 420  
|||||  
Db 61 lghpkrkaritthcngmgiyeadityhgvpmvvpifgpolninahmkakaveinfkrtmts 120

OY 421 EDLRLRLT 429  
|||  
Db 121 edledalks 129

#### RESULT 15

AA24411  
ID AAB44411 standard; Protein; 129 AA.

XX AAB44411;

XX 05-FEB-2001 (first entry)

XX Human lung tumour-specific antigen encoded by cDNA #21.

XX Lung tumour protein; lung cancer; cytostatic; vaccine.

XX Homo sapiens.

XX WO200060077-A2.

XX 12-OCT-2000.

XX 30-MAR-2000; 2000MO-US08560.

XX 02-APR-1999; 99US-0285323.

XX 09-AUG-1999; 99US-0370838.

XX 30-DEC-1999; 99US-0476235.

XX 03-MAR-2000; 2000US-0518809.

XX (CORI-) CORIXA CORP.

XX Reed SG, Lodes MJ, Mohamath R, Secrist H;

XX WPI; 2000-638466/61.

XX N-PSDB; AAC79066.

XX Novel lung tumour polypeptides and polynucleotides, useful for  
XX detecting, monitoring or treating cancer, especially lung cancer -

XX Claim 1; Page 99; 243pp; English.

XX The present sequence is given in a specification relating to compounds  
XX for therapy and diagnosis of lung cancer. Polypeptides comprising at  
XX least an immunogenic part of a lung tumour protein are disclosed.

XX The polypeptides are useful for inhibiting the development of cancer,  
XX especially lung cancer. Samples of T cells expressing the polypeptides  
XX may be used to inhibit the development of cancer. The polypeptides are  
XX also useful for detecting and monitoring the progression of cancer,  
XX especially lung cancer.

XX Sequence 129 AA;

